

10. Rollino C, Garofalo G, Roccatello D *et al.* Colour-coded Doppler sonography in monitoring native kidney biopsies. *Nephrol Dial Transplant* 1994; 9: 1260–1263
11. Christensen J, Lindequist S, Knudsen DU *et al.* Ultrasound-guided renal biopsy with biopsy gun technique—efficacy and complications. *Acta Radiol* 1995; 36: 276–279
12. Diaz-Buxo JA, Donadio JV Jr. Complications of percutaneous renal biopsy: an analysis of 1000 consecutive biopsies. *Clin Nephrol* 1975; 4: 223–227
13. Stratta P, Canavese C, Marengo M *et al.* Risk management of renal biopsy: 1387 cases over 30 years in a single centre. *Eur J Clin Invest* 2007; 37: 954–963
14. Shidham GB, Siddiqi N, Beres JA *et al.* Clinical risk factors associated with bleeding after native kidney biopsy. *Nephrology (Carlton)* 2005; 10: 305–310
15. Birnholz JC, Kasinath BS, Corwin HL. An improved technique for ultrasound guided percutaneous renal biopsy. *Kidney Int* 1985; 27: 80–82
16. Bach D, Wirth C, Schott G *et al.* Percutaneous renal biopsy: three years of experience with the biopsy gun in 761 cases—a survey of results and complications. *Int Urol Nephrol* 1999; 31: 15–22
17. Fraser IR, Fairley KF. Renal biopsy as an outpatient procedure. *Am J Kidney Dis* 1995; 25: 876–878
18. Nass K, O'Neill WC. Bedside renal biopsy: ultrasound guidance by the nephrologist. *Am J Kidney Dis* 1999; 34: 955–959
19. Nyman RS, Cappelen-Smith J, al Suhaibani H *et al.* Yield and complications in percutaneous renal biopsy. A comparison between ultrasound-guided gun-biopsy and manual techniques in native and transplant kidneys. *Acta Radiol* 1997; 38: 431–436
20. Lin WC, Yang Y, Wen YK *et al.* Outpatient versus inpatient renal biopsy: a retrospective study. *Clin Nephrol* 2006; 66: 17–24
21. Health and Public Policy Committee, American College of Physicians. Clinical competence in percutaneous renal biopsy. *Ann Int Med* 1988; 108: 301–303
22. Mattix H, Singh AK. Is the bleeding time predictive of bleeding prior to a percutaneous renal biopsy? *Curr Opin Nephrol Hypertens* 1999; 8: 715–718
23. Peterson P, Hayes TE, Arkin CF *et al.* The preoperative bleeding time test lacks clinical benefit. *Arch Surg* 1998; 133: 134–139
24. Bolton WK. Nonhemorrhagic decrements in hematocrit values after percutaneous renal biopsy. *JAMA* 1977; 238: 1266–1268
25. Bolton WK, Gibson RS, Eells PF. Vasovagal pseudo-hemorrhage. Complication of percutaneous renal biopsy. *JAMA* 1977; 237: 1259–1260
26. Burstein DM, Schwartz MM, Korbet SM. Percutaneous renal biopsy with the use of real-time ultrasound. *Am J Nephrol* 1991; 11: 195–200
27. Khajehdehi P, Junaid SM, Salinas-Madriral L *et al.* Percutaneous renal biopsy in the 1990s: safety, value, and implications for early hospital discharge. *Am J Kidney Dis* 1999; 34: 92–97
28. Ralls PW, Barakos JA, Kaptein EM *et al.* Renal biopsy related hemorrhage: frequency and comparison of CT and sonography. *J Comput Assist Tomogr* 1987; 11: 1031–1034
29. Rosenbaum R, Hoffsten PE, Stanley RJ *et al.* Use of computerized tomography to diagnose complications of percutaneous renal biopsy. *Kidney Int* 1978; 14: 87–92
30. Ginsburg JC, Fransman SL, Singer MA *et al.* Use of computerized tomography to evaluate bleeding after renal biopsy. *Nephron* 1980; 26: 240–243
31. Riehl J, Maigatter S, Kierdorf H *et al.* Percutaneous renal biopsy: comparison of manual and automated puncture techniques with native and transplanted kidneys. *Nephrol Dial Transplant* 1994; 9: 1568–1574
32. Doyle AJ, Gregory MC, Terreros DA. Percutaneous native renal biopsy: comparison of a 1.2 mm spring-loaded system with a traditional 2 mm hand-driven system. *Am J Kidney Dis* 1994; 23: 498–503

Received for publication: 20.11.08; Accepted in revised form: 2.2.09

Nephrol Dial Transplant (2009) 24: 2439–2445

doi: 10.1093/ndt/gfp101

Advance Access publication 17 March 2009

The correlation between dental calculus and disturbed mineral metabolism in paediatric patients with chronic kidney disease

Esti Davidovich^{1,*}, Miriam Davidovits^{2,3,*}, Benny Peretz⁴, Joseph Shapira¹ and Doron J. Aframian⁵

¹Department of Pediatric Dentistry, Hebrew University, Hadassah Faculty of Dental Medicine, Jerusalem, ²Institute of Paediatric Nephrology, Schneider Children's Medical Center of Israel, Petah Tiqva, Israel, ³Sackler School of Medicine, Tel Aviv University, ⁴Department of Pediatric Dentistry, The Maurice and Gabriela Goldschleger School of Dental Medicine, Tel Aviv University, Tel Aviv and ⁵Department of Oral Medicine, Salivary Gland Clinic, The Hebrew University-Hadassah School of Dental Medicine, Jerusalem, Israel

Correspondence and offprint requests to: Doron J. Aframian; E-mail: dorona@cc.huji.ac.il

*Both these authors contributed equally to the study.

Abstract

Background. Vascular calcifications have been documented in children with end-stage renal disease. However, only a few reports have described abundant dental calculus formation in children suffering from chronic kidney dis-

ease (CKD). Moreover, dental calculus scores (DCS) and their correlation with renal disease severity have not been studied.

Methods. DCS in 74 young CKD patients were evaluated: 25 pre-dialytic (PrD), 18 on dialysis (D) and 31 with

transplants (T) compared to 32 healthy participants (C). Saliva and serum analysis included creatinine (Cr), urea (U), calcium (Ca), phosphorous (P), magnesium (Mg) as well as intraoral pH levels.

Results. All patient groups presented high DCS. DCS and pH levels were higher in the D group with a positive correlation between pH and lower incisor DCS ($r = 0.56$, $P = 0.017$). The highest salivary Ca was found in the PrD group. Salivary P in the PrD group was found to be higher than in the T and C groups. The lowest salivary Mg was found in the D group while the highest salivary Ca $\times$ P product was found in the PrD group. In all patient groups, salivary U was higher than in the C group with a 2.5-fold increase in the D group. Salivary Cr resembled the U salivary concentrations.

Conclusions. Alterations in salivary Ca, P, Mg, U, Cr and intraoral pH levels were observed in the patient groups. DCS correlated with renal disease severity and therefore may be a reflection of other tissue calcification pathologies found in these patients.

Keywords: Ca $\times$ P product; chronic kidney disease; dental calculus; saliva; vascular calcifications

Introduction

Dental calculus in healthy children is rare [1,2]; however, several studies have reported an elevated amount of calculus in the paediatric population with chronic kidney disease (CKD) [3–5].

Recently, we have reported a case with an abundant amount of dental calculus in an infant on dialysis that totally disappeared after kidney transplantation [6].

Dental calculus is a form of a calcification process in the oral cavity environment where calcium (Ca) and phosphorous (P) ions originating from saliva play a major role. This calculus is assembled through interaction with the dental plaque, which is a community of microorganisms found on the tooth surface as a biofilm [7]. In a healthy oral environment, saliva is supersaturated with Ca and P levels, however, without precipitation. Nonetheless, when this equilibrium is disturbed, dental calculus is formed enhanced by elevated salivary pH [8]. Dental calculus is formed with four different crystals of Ca–P; brushite, octa Ca–P, hydroxyapatite and whitlockite in which the most abundant crystals are the hydroxyapatite and octa Ca–P [9].

CKD is characterized by profound disturbances in Ca–P metabolism and increased Ca burden as a result of administration of Ca-based phosphate binders [10–14]. Cardiovascular complications related to abnormal Ca–P metabolism are the major cause of morbidity and mortality observed in CKD patients and particularly in end-stage renal disease (ESRD) [10–13]. Moreover, different manifestations of uraemic vasculopathy due to calcifications have been reported recently among the paediatric CKD population [14–19].

Since the quantification of dental calculus and its correlation with renal disease severity have not been previously studied in young CKD patients, our objectives were (1) to evaluate the dental calculus formation in young CKD

Table 1. Clinical data of 106 participants in the study

Variables	Control	Pre-dialysis	Dialysis	Transplanted
Gender (M/F)	23/9	14/11	11/7	20/11
Age (years) $\pm$ SD	9.2 $\pm$ 2.7	12.6 $\pm$ 5	14.4 $\pm$ 3	14.7 $\pm$ 5
GFR (mean $\pm$ SEM, range)	ND	40.44 $\pm$ 3.72, 13.7–67.3*	NR	58.61 $\pm$ 3.15, 11.59–95.9**

GFR = glomerular filtration rate, ND = not determined, NR = not relevant.

*Significant difference between pre-dialytic and transplant groups.

**Significant difference from transplant group.

patients and (2) to examine salivary mineral metabolism and intraoral mucosal pH levels with correlation to dental calculus formation in different stages of kidney function impairment.

Methods

Study population

Upon approval by the Institutional Review Board for research on human subjects and informed consent, 106 children, adolescents and young adults were recruited and studied.

The CKD patients were divided as follows: 25 pre-dialytic (PrD) patients, 18 patients on maintenance dialysis (8 on haemodialysis, 10 on peritoneal dialysis) (group D), and 31 transplanted patients with a functioning graft (group T).

All patients were treated at the Institute of Paediatric Nephrology, Schneider Children's Medical Center of Israel, Petah Tiqva, Israel.

Thirty-two healthy children without previous history of impaired renal function served as the control group (group C). This group was recruited and examined in the Department of Pediatric Dentistry at the Hebrew University—Hadassah School of Dental Medicine, Jerusalem, Israel (Table 1).

Systemic variables

The glomerular filtration rate (GFR) was calculated from the serum creatinine (Cr) according to the Schwartz formula [20]. Serum Cr, urea (U), Ca, P and magnesium (Mg) concentrations were measured using the Hitachi 917 autoanalyser (Roche Ltd, Basel, Switzerland).

Oral measurements

All intraoral assessments were utilized by one examiner (ED). The degree of dental calculus formation was scored according to WHO oral hygiene indices [21,22]. Briefly, a score of 0 represents no calculus formation whereas a score of 3 represents supragingival calculus covering more than two-thirds of the exposed tooth surface or a continuous heavy band of subgingival calculus around the cervical portion of the tooth or both. Two locations were examined: (1) the lingual aspects of the mandibular incisors adjacent to the orifices of the submandibular glands and (2) the buccal aspects of the right and left first upper molars adjacent to the orifices of the parotid glands.

Unstimulated whole saliva was collected using the spitting method as previously described [23–26]. In brief, saliva collection was done in a quiet room between 8:00 and 12:00 AM. The subjects were refrained from eating, tooth brushing or using mouth wash rinses at least 1 h before spitting. They were asked to collect saliva in their mouth and to spit into a wide test tube for 5 min. Thereafter, the collected saliva was immediately delivered at 4°C and further kept at –80°C until analysis was performed.

Saliva samples were thawed, and then centrifuged (2000 g, 20 min, 250°C). The supernatants were analysed using the Hitachi 917 autoanalyser (Roche Ltd), and the concentrations of U, Cr, Ca, P and Mg were measured. Measurement of citrate (Cit) concentrations was performed in five healthy subjects (C group) and five dialysis patients (D group) using the UV method with the Boehringer Mannheim kit (Darmstadt, Germany).

Table 2. Dental calculus score for all groups

DSC (mean $\pm$ SD)	Control	Pre-dialysis	Dialysis	Transplanted
Lower incisors	0	1.92 $\pm$ 1.25*•	3.44 $\pm$ 1.04*	1.29 $\pm$ 1.16*•
Upper molars	0	1.28 $\pm$ 1.06*•	2.67 $\pm$ 1.14*	0.77 $\pm$ 0.8*•

*Significant difference from the control group.

•Significant difference from the dialysis group.

The method is based on converting citric acid to oxaloacetate and acetate (by the enzyme citrate lyase), and then by other two enzymes (L-malate dehydrogenase and L-lactate dehydrogenase) reducing NADH to NAD+ that is stoichiometric to the amount of citrate. Reference intervals for 24-h urine collection are 290–1150 mg/24 h. The analytical sensitivity is 0.25 mg/dl.

Oxalate (Ox) levels in saliva were utilized in five healthy subjects (C group) and five dialysis patients (D group) with the Trinity Biotech kit (Wicklow, Ireland). Ox analysis in saliva was based on urine analysis and was performed on Olympus 2700 analyser using an enzymatic method. The method is based on converting oxalate to CO₂ and H₂O₂ (by the enzyme oxalate oxidase), followed by H₂O₂ reaction with 3-methyl-2-benzothiazolinone hydrazone (MBTH) and 3-(dimethylamino) benzoic acid (DMAB) in the presence of peroxidase, to yield an indamine dye. The intensity of the colour produced is directly proportional to the concentration of Ox in the sample. Reference intervals for 24-h urine collection are 7–45 mg/24 h for male, 4–31 mg/24 h for female and 13–38 mg/dl for children and were implemented to saliva calculation. The analytical sensitivity was 0.88 mg/dl.

Calculus composition was assessed in five dialysis patients using urinary calculi analysis kit (DiaSys Diagnostic Systems GmbH, Holzheim, Germany).

The oral surface pH was assessed with a flat, glass electrode pH meter (HANNA instruments HI 8424, Padova, Italy) as previously described [26]. Two sets of measurements were applied at seven locations: the soft and hard palate, anterior, middle, and posterior tongue, right and left buccal mucosa. Five minutes were allowed between measurements [26].

Statistical analysis

Frequencies and percentages were calculated for categorical variables. The frequencies between categorical variables were analysed by the 'chi square test' and the 'Fisher-Irwin exact test'. Medians, means and standard errors were calculated for continuous parameters. The results of pH parameters between subgroups of patients were compared by 'one-way analysis-of-variance' using the Bonferroni model. Due to large variability of salivary components, the 'Kruskal-Wallis' test was used to compare between saliva parameters. The results between pairs of saliva parameters were analysed using the Wilcoxon signed-rank test. The correlations between DCS and pH levels were analysed by 'Pearson's correlation coefficient' and the correlations between saliva parameters were analysed by 'Spearman's correlation coefficient'. The significance level of the correlation coefficient was calculated.

Results

Dental calculus score

Calculus was not found in any individual of the C group (Table 2) as opposed to calculus formation in all the patient groups. The highest DCS was found in the D group with a significant difference from the PrD and T groups in both upper and lower sites ($P = 0.0001$). The T group patients exhibited the lowest DCS; however, with no significant difference from the PrD group. The scores were higher at the lower incisor site compared to the molar site in all study groups (Table 2).

Serum and saliva parameters

The serum parameters including Cr, Ca, P, Mg and U (group D only) are listed in Table 3.

The salivary Ca level in the PrD group was significantly higher compared to the D, C and T groups (3.7-fold $P = 0.0001$, 2.5-fold $P = 0.05$, 1.5-fold $P = 0.001$, respectively). The Ca levels in the T group were found to be significantly higher than in the D group (means of 2.96 mg/dl and 1.18 mg/dl, respectively, $P = 0.0001$) and in the C group (means of 2.96 mg/dl and 1.78 mg/dl, respectively, $P = 0.001$). (Table 3).

Salivary P levels in the PrD group (mean = 21.5 mg/dl) were found to be significantly higher than in the T group (mean 15.4 mg/dl, $P = 0.015$) and C groups (mean 14.35 mg/dl, $P = 0.001$). In the D group, P levels were found to be higher than in the C group although not statistically significant (Table 3).

The salivary Mg concentrations in the D group (mean = 0.25 mg/dl) were found to be significantly lower than in the PrD (mean = 0.55 mg/dl, $P = 0.0004$), T (mean = 0.57 mg/dl, $P = 0.002$) and C groups (mean = 0.43 mg/dl, $P = 0.017$). The Mg values in the PrD group were found to be significantly higher than in the C group ($P = 0.048$) as depicted in Table 3.

The salivary Ca $\times$ P product in the PrD group was significantly higher (mean 117.74) than in the D, T and C groups ($P = 0.0002$, $P = 0.01$, $P = 0.0001$, respectively). The Ca $\times$ P product for the T group was also significantly higher than in the C and D groups ($P = 0.002$, $P = 0.01$, respectively) as shown in Table 3.

In all the patient groups D, PrD and T (mean = 83.9, 80.5, 52.7 mg/dl, respectively), salivary U concentration was significantly higher than in the C group (mean 31.6 mg/dl; $P = 0.002$, $P = 0.0001$, $P = 0.0001$, respectively) with more than a 2.5-fold increase in the D group followed by the PrD and the T groups. A significant difference was also found between the PrD group and the T group ($P = 0.007$) (Table 3).

A significant correlation was found between GFR and salivary U concentrations in the PrD and T groups ($r = -0.66$, $P = 0.001$, $r = -0.45$, $P = 0.022$, respectively). Also, a significantly higher salivary U concentration was found in the D group followed by the PrD and the T groups (Table 3) and corresponding to DCS (Table 2).

The Cr salivary concentration pattern resembled the urea salivary concentration with significantly higher levels in the D group (mean 0.59 mg/dl) followed by the PrD (mean 0.4 mg/dl), T (mean 0.15 mg/dl) and C groups (mean 0.14 mg/dl) (Table 3).

Oral mucosal pH scores

Table 4 represents the average of pH scores from seven intraoral mucosal sites. The average of pH values in the D group was significantly higher (mean 7.38) than in the PrD group (mean 6.82, $P = 0.001$), the T group (mean 6.73, $P = 0.0001$) and the C group (mean 6.72, $P = 0.0001$). No statistically significant difference was found in pH scores between the PrD, the T and the C groups. A correlation between the average pH score and the DCS at the lower incisor

Table 3. Serum and salivary biochemistry parameters in all examined groups

Parameter [mean $\pm$ SEM (range) median]	Control		Pre-dialysis		Dialysis		Transplanted	
	Blood ^a	Saliva	Blood	Saliva	Blood	Saliva	Blood	Saliva
Urea (mg/dl)	2–50	31.6 $\pm$ 2.8*, [^] (14–76) 26.5	ND	80.5 $\pm$ 8.3*, [^] , [×] (16–142) 77	122.7 $\pm$ 8.4 (74–210) 115.5	83.9 $\pm$ 16.2* (16–197) 78.5	ND	52.7 $\pm$ 5.0*, [×] (23–141) 41
Creatinine (mg/dl)	0.5–1.1	0.14 $\pm$ 0.02*, [×] (0.06–0.44) 0.11	2.06 $\pm$ 0.27 (0.81–5.7) 1.5	0.40 $\pm$ 0.08*, [^] (0.02–2.1) 0.31	7.28 $\pm$ 0.36 (4.48–9.7) 7.15	0.59 $\pm$ 0.1* [^] (0.13–1.47) 0.51	1.28 $\pm$ 0.11 (0.66–3.96) 1.17	0.15 $\pm$ 0.02*, [×] (0.02–0.58) 0.11
Calcium (mg/dl)	9.2–11	1.79 $\pm$ 0.17 [^] , [×] (0.5–4) 1.55	9.78 $\pm$ 0.2 (6.9–11.5) 10	4.41 $\pm$ 0.59*, [•] , [^] (1.5–11.1) 3.42	9.5 $\pm$ 0.2, (8.4–11.4) 9	1.18 $\pm$ 0.22 [^] , [×] (0.2–2.2) 1.2	10.0 $\pm$ 0.08 (9.2–11.2) 10	2.96 $\pm$ 0.28*, [•] , [×] (0.9–6.3) 2.8
Phosphorus (mg/dl)	3–6	15.82 $\pm$ 1.15 [×] (7.2–41.2) 14.35	5.0 $\pm$ 0.25 (2.7–9.4) 4.6	23.22 $\pm$ 2.10 [^] , [*] (9.3–52.3) 21.5	5.72 $\pm$ 0.27 (4.1–8.9) 5.4	20.77 $\pm$ 2.57 (7–36) 19.55	5.8 $\pm$ 1.4, (2.6–47.0) 4.5	17.44 $\pm$ 1.46 [×] (7.7–42.5) 15.4
Magnesium (mg/dl)	1.6–2.4	0.46 $\pm$ 0.05*, [×] (0.12–1.1) 0.43	ND	0.81 $\pm$ 0.16*, [•] (0.15–3.45) 0.55	ND	0.25 $\pm$ 0.05*, [^] , [×] (0.01–0.47) 0.25	ND	0.64 $\pm$ 0.07*, [•] (0.17–1.67) 0.57
Calcium $\times$ phosphorus product	30–55	27.9 $\pm$ 3.5*, [^] , [×] (7.35–107.1) 22.75	48.45 $\pm$ 2.32 (24.84–80.84)	117.74 $\pm$ 28.07*, [•] , [^] (55.86–528.13) 67.75	54.65 $\pm$ 3.07 (36.9–89.89)	25.56 $\pm$ 5.92 [^] , [×] (2.8–67.98) 24.66	59.4 $\pm$ 14.7 (26.78–498.2)	52.96 $\pm$ 7.05*, [•] , [×] (11.7–161.55) 44.52

ND = not determined.

^aNormal serum range adopted from [38].

*Significant difference from the control group.

•Significant difference from the dialysis group.

^Significant difference from the pre-dialysis group.

×Significant difference from the transplant group.

Table 4. Intraoral mucosal pH score

pH (mean $\pm$ SD, range)	Control	Pre-dialysis	Dialysis	Transplanted
All sites	6.72 $\pm$ 0.06, 5.84–7.38**	6.82 $\pm$ 0.12, 5.60–7.85**	7.38 $\pm$ 0.13, 5.95–8.03	6.73 $\pm$ 0.07, 5.90–7.70**
Hard palate	7.18 $\pm$ 0.13, 5.92–9.18*	7.05 $\pm$ 0.15, 5.77–8.45**	7.73 $\pm$ 0.14, 6.46–9.6	6.95 $\pm$ 0.09, 6.05–8.24**
Soft palate	7.08 $\pm$ 0.12, 5.53–8.2*	7.1 $\pm$ 0.15, 5.74–8.47*	7.7 $\pm$ 0.16, 6.16–8.54	6.95 $\pm$ 0.1, 6.08–8.5**
Right mucosa	6.46 $\pm$ 0.06, 5.75–7.42**	6.58 $\pm$ 0.1, 5.39–7.88**	7.05 $\pm$ 0.13, 5.92–7.75	6.42 $\pm$ 0.06, 5.5–6.88**
Left mucosa	6.4 $\pm$ 0.07, 5.4–7.43**	6.56 $\pm$ 0.1, 5.5–7.58*	7.00 $\pm$ 0.13, 5.78–7.71	6.44 $\pm$ 0.07, 5.5–7.65**
Posterior tongue	6.58 $\pm$ 0.09, 5.92–9.18**	6.80 $\pm$ 0.15, 5.36–7.95**	7.41 $\pm$ 0.16, 5.64–8.51	6.77 $\pm$ 0.08, 5.9–7.51**
Middle tongue	6.63 $\pm$ 0.07, 5.72–7.35**	6.83 $\pm$ 0.14, 5.35–7.78**	7.45 $\pm$ 0.15, 5.82–8.51	6.82 $\pm$ 0.07, 5.96–7.6**
Anterior tongue	6.71 $\pm$ 0.06, 5.84–7.38**	6.82 $\pm$ 0.12, 5.6–7.85**	7.35 $\pm$ 0.13, 5.90–8.37	6.77 $\pm$ 0.06, 6.15–7.58**

Significant difference from the control group (* $P \leq 0.05$; ** $P \leq 0.01$).

*Significant difference from the dialysis group.

× Significant difference from the pre-dialysis group.

^Significant difference from the transplant group.

site was found in the D group (correlation coefficient = 0.56, $P = 0.017$).

Calculus composition and salivary Ox and Cit concentrations in saliva

Table 5 represents the Ca, P and Ox levels obtained from dental calculus in five D patients. High levels of Ox were found followed by P and Ca (means 19, 14 and 12%, respectively) levels. Ox and Cit were measured in saliva of five dialysis patients and five healthy age-matched children (Table 5). More than a 3-fold increase in Ox levels was found in the D group with only slight increase in the Cit levels compared to the C group.

Discussion

To our knowledge, this study is the first to examine the dental calculus formation in young patients with different stages of chronic kidney disease (CKD) including pre-dialytic (PrD) patients on dialysis (D) and patients after kidney transplantation (T) compared to control healthy children. Considering the mechanism of the calcification process and the metabolic disturbances resulting from kidney dysfunction, we explored the Ca, P, U, Cr and Mg concentrations in the secreted saliva of these patients and the pH levels in intraoral mucosal sites.

DCS were found to be higher in the lingual aspect of the lower incisors in all the study groups (Table 2). This finding is consistent with previous observations of preferential calculus formation on the lingual surface of the six lower incisors due to their proximity to the orifices of the submandibular glands, which act as the reservoir of Ca and P ions [8]. Interestingly, when we further explored the calculus composition in the most affected D patients (Table 5), we found that the Ox component was observed. To our knowledge, no previous studies have explored Ca–Ox as part of dental calculus specifically in CKD patients. The fact that an elevated concentration of Ox in saliva was observed is probably due to the decrease in GFR and urinary excretion in these patients. Higher oral pH levels were established in the D group patients versus the C group (pH = 7.38, pH =




Fig. 1. Correlation between salivary parameters and dental calculus score in chronic renal failure disease stage. DCS = dental calculus score, CRF = chronic renal failure, PrD = pre-dialysis group, D = dialysis group, T = transplant group, ↑ = high salivary concentration, ↓ = low salivary concentration, U = urea, Ca = calcium, Mg = magnesium.

6.72, respectively, Table 3). In this condition, Ca–Ox saturation product becomes extremely high due to an equimolar increase in Ox as compared to Ca. Consequently, it will lead to a 10-fold higher increase in the saturation process and hence to higher risk in crystal deposition.

Dental calculus was not found in control subjects (Table 2) similar to previous studies [1,2]. Interestingly, the most abundant amount of calculus was found in the D patient group, followed by the PrD group and the T group, clearly demonstrating that deterioration of the renal function is accompanied by calculus formation reaching its peak incidence in ESRD, and significantly improved after kidney transplantation.

Blood U and Cr levels serve as indicators for the severity of kidney disease. Previous studies have shown elevated levels of salivary U concentrations in patients with ESRD [27,28]. These observations correlate with our study where both salivary U and Cr levels changed in saliva according to the renal function, with the highest levels found in the D group (Table 3). U has been shown to facilitate dental plaque alkalization [8], and therefore may contribute to the highest rate of calculus formation in the D group (Figure 1). In the PrD and T groups, a significant

Table 5. Calculus composition in dialysis patients and salivary oxalate and citrate concentrations in dialysis group versus controls

	Calculus composition (%)			Saliva composition (mg/dl)			
	D group			D group		C group	
#	Ca	Ox	P	Ox	Cit	Ox	Cit
1	5	10	5	0.12	8.1	0.03	0.6
2	10	30	15	0.07	17.8	ND	8.3
3	15	25	15	0.12	7.2	0.04	1.9
4	25	10	20	ND	7.2	ND	17.1
5	5	20	15	0.08	8.7	0.02	14
Mean $\pm$ SD	12 $\pm$ 8.4	19 $\pm$ 8.9	14 $\pm$ 5.5	0.098 $\pm$ 0.05	9.8 $\pm$ 4.5	0.03 $\pm$ 0.02	8.4 $\pm$ 7.3

Calcium, oxalate and phosphate composition was obtained from calculus. Oxalate and citrate were measured in saliva of five dialysis patients and five healthy age-matched children. Ca = calcium; P = phosphorus; Ox = oxalate; Cit = citrate; ND = not detected; D = dialysis group; C = control group.

correlation was found between the GFR and the salivary U concentrations (results are not shown). This observation may help in the future to assess renal function in these two groups (i.e. PrD and T) by measuring salivary U concentration.

The highest measurement of oral mucosal pH was found in the D group corresponding to the salivary U concentrations in this group. Moreover, a correlation between average pH measurements and lower incisor DCS was found in this group (correlation coefficient = 0.56, P = 0.017). This observation can be explained by the abundant supply of U from the salivary gland secretion hydrolysed by bacteria with the release of ammonia, leading to an elevation of pH in the dental plaque and further promoting Ca and P precipitation [29,8].

The most significantly higher concentration of salivary Ca was found in the PrD group. This finding may be a reflection of the disturbed Ca metabolism in this patients' group manifested by the high Ca burden followed by compensatory secretion in saliva. Salivary Ca levels during dialysis were reduced, probably in parallel to the reduction of Ca levels in the serum at this stage. A further decrease in salivary Ca secretion in the T group may be attributed to a decrease in the Ca burden due to the improved renal function and cessation of Ca-containing P-binder medications. Interestingly, Ca levels were still higher in the T group compared to controls, which is probably related to the still existing Ca overload state (Table 2).

P concentrations in the saliva in all groups were higher than in the serum reaching almost a 5-fold increase in the PrD group corresponding to the active mechanism of P secretion from the salivary glands [30]. Recently, Savica *et al.* [31] have found a significant increase in salivary P levels in adult haemodialysis patients compared to healthy controls; serum phosphorus was the only predictor of increased excretion in saliva. The authors suggested that increased salivary P secretion might be interpreted as being compensatory in the presence of renal failure [31].

Consequently, upon renal function deterioration, more P is secreted in the saliva leading to precipitation with Ca and dental calculus formation as observed in this study (Figure 1).

Elevated plasma P levels and high Ca $\times$ P ion product play a key role in vascular calcifications, increasing the

prevalence of morbidity and mortality in CKD patients [13,16,17,32–34]. Furthermore, several studies have shown that long-term exposure to disturbed mineral metabolism in paediatric CKD patients appears to contribute to the development of coronary artery calcifications in young adults with ESRD [13,19,35,36]. Vascular abnormalities are already present in children and adolescents during early stages of CKD. These abnormalities are more severe in children on dialysis, improve after kidney transplantation and are related to abnormal Ca–P metabolism. In recent years, attention has focused on the mechanism of the calcification process, its detection and monitoring as well as the role of Ca–P as potentially modifiable risk factor in the development and progression of vascular calcifications [16–18]. In this study, the salivary Ca $\times$ P product was significantly higher in the PrD group followed by the T group (Table 3). Interestingly, Ca $\times$ P product levels in the serum did not significantly change between the groups. This discrepancy between serum and saliva may be attributed to higher salivary P secretion as previously discussed, emphasizing the key role of P in dental calculus formation.

The lowest salivary Mg concentration was found in the D group (Table 1). Mg acts as an inhibitor ion of the calcification process further explaining the amplification of dental calculus formation in the D group. Interestingly, Grases *et al.* [37] showed previously less sialolithiasis formation in patients with higher salivary Mg concentrations [36].

In conclusion, our study clearly demonstrates a possible association between the severity of renal dysfunction in young patients and the formation of dental calculus—an additional manifestation of disturbed Ca–P homeostasis.

The combination of several components in saliva including Ca, P, U and Mg plays an important role in this process. The elevated Ox content finding in the D group patients' saliva as well as a major component in their dental calculi may also contribute to the understanding of calculus formation process in CKD patients.

Acknowledgements. This study was sponsored by the Israeli Ministry of Health, grant no. 3-6396. We thank Mrs Shlomit Gan for her statistical work for this manuscript and Dr Ruth Tor for her assistance. Partial data from this manuscript were presented as an abstract in the IADR 2007 Thessaloniki conference.

Conflict of interest statement. None declared.

References

- McDonald RE, Avery DR, Weddel JA. Gingivitis and periodontal diseases. In: McDonald RE, Avery DR, Dean JA, (eds). *Dentistry for the Child and Adolescent*. St Louis, MO: Mosby, 2004, 449–450
- Baht M. Periodontal health of 14–17-year-old US school children. *J Public Health Dent* 1991; 51: 5–11
- Epstein S, Mandel I, Scopp I. Salivary composition and calculus formation in patients undergoing hemodialysis. *J Periodontol* 1980; 51: 336–338
- Obry F, Belcourt AB, Frank RM *et al.* Biochemical study of whole saliva from children with chronic renal failure. *J Den Child* 1987; 54: 429–432
- Martins C, Siqueira WL, de Oliveira E *et al.* Salivary analysis of patients with chronic renal failure undergoing hemodialysis. *Spec Care Dentist* 2006; 26: 205–208
- Davidovich E, Frishberg Y, Aframian DJ *et al.* Calculus in a toddler with end-stage renal disease due to prune-belly syndrome. *Oral Dis* 2006; 12: 63–66
- Marsh PD. Dental plaque as a microbial biofilm. *Caries Res* 2004; 38: 204–211
- Dawes C. Why does supragingival calculus form preferentially on the lingual surface of the 6 lower anterior teeth? *J Can Dent Assoc* 2006; 72: 923–926
- Lang NP, Mombelli R, Attstrom R. Oral biofilms and calculus. In: Linde J, Lang NP, Karring T. (eds). *Clinical Periodontology and Implant Dentistry*. Oxford, UK: Blackwell Munksgaard, 2008, 197–205
- Blacher J, Pannier B, Guerin AP *et al.* Carotid arterial stiffness as a predictor of cardiovascular and all-cause mortality in end-stage renal disease. *Hypertension* 1998; 32: 570–574
- Raggi P, Boulay A, Chasan-Taber S *et al.* Cardiac calcification in adult hemodialysis patients. A link between end-stage renal disease and cardiovascular disease? *J Am Coll Cardiol* 2002; 39: 695–701
- Groothoff JW, Gruppen MP, Offringa M *et al.* Mortality and causes of death of end-stage renal disease in children: a Dutch cohort study. *Kidney Int* 2002; 61: 621–629
- Goodman W, Goldin J, Kuizon B *et al.* Coronary-artery calcification in young adults with end-stage renal disease who are undergoing dialysis. *N Engl J Med* 2000; 342: 1478–1483
- Civilibal M, Caliskan S, Adaletli I *et al.* Coronary artery calcifications in children with end-stage renal disease. *Pediatr Nephrol* 2006; 21: 1426–1433
- Covic A, Mardare N, Gusbeth-Tatomir P *et al.* Increased arterial stiffness in children on haemodialysis. *Nephrol Dial Transplant* 2006; 21: 729–735
- Litwin M, Wühl E, Jourdan C *et al.* Altered morphologic properties of large arteries in children with chronic renal failure and after renal transplantation. *J Am Soc Nephrol* 2005; 16: 1494–1500
- Mitsnefes MM, Kimball TR, Kartal J *et al.* Cardiac and vascular adaptation in pediatric patients with chronic kidney disease: role of calcium-phosphorus metabolism. *J Am Soc Nephrol* 2005; 16: 2796–2803
- Shrorff RC, Donald AE, Hiorns MP *et al.* Mineral metabolism and vascular damage in children on dialysis. *J Am Soc Nephrol* 2007; 18: 2996–3003
- Oh J, Wunsch R, Turzer M *et al.* Advanced coronary and carotid arteriopathy in young adults with childhood-onset chronic renal failure. *Circulation* 2002; 106: 100–105
- Schwartz GJ, Haycock GB, Edelman CM Jr. A simple estimate of glomerular filtration rate in children derived from body length and plasma creatinine. *Pediatrics* 1976; 58: 259–263
- WHO oral hygiene indices <http://www.whocollab.od.mah.se/exp/ohigv60.html>
- Conroy C, Sturzenberger O. The rate of calculus formation in adults. *Periodontol* 1968; 39: 142–144
- Ben-Aryeh H, Shalev A, Szargel R *et al.* The salivary flow rate and composition of whole and parotid resting and stimulated saliva in young and old healthy subjects. *Biochem Med Metab Biol* 1986; 36: 260–265
- Sagulin G, Roomans G. Effects of thyroxine and dexamethasone on rat submandibular glands. *J Dent Res* 1998; 68: 1247–1251
- Melvin J. Saliva and dental diseases. *Curr Opin Dent* 1991; 1: 795–801
- Aframian DJ, Davidowitz T, Benoliel R. The distribution of oral mucosal pH values in healthy saliva secretors. *Oral Dis* 2006; 12: 420–423
- Peterson S, Woodhead J, Crall J. Caries resistance in children with chronic renal failure: plaque pH, salivary pH, and salivary composition. *Pediatr Res* 1985; 19: 796–799
- Al-Nowaiser A, Roberts GJ, Trompeter RS. Oral health in children with chronic renal failure. *Pediatr Nephrol* 2003; 18: 39–45
- Sissons CH, Hancock EM, Cutress TW. The sources of variation in ureolysis in artificial plaque cultured from human salivary bacteria. *Arch Oral Biol* 1988; 33: 721–726
- Rehak NN, Cecco SA, Csako G. Biochemical composition and electrolyte balance of ‘unstimulated’ whole human saliva. *Clin Chem Lab Med* 2000; 38: 335–343
- Savica V, Calo LA, Caldarella R *et al.* Phosphatase salivary secretion in hemodialysis patients: implications for the treatment of hyperphosphatemia. *Nephron Physiol* 2007; 105: 52–55
- Block GA. Prevalence and clinical consequences of elevated Ca × P product in hemodialysis patients. *Clin Nephrol* 2000; 54: 318–324
- Ganesh SK, Stack AG, Levin NW *et al.* Association of elevated serum PO(4), Ca × PO(4) product, and parathyroid hormone with cardiac mortality risk in chronic hemodialysis patients. *J Am Soc Nephrol* 2001; 12: 2131–2138
- Giachelli CM. Vascular calcification mechanisms. *J Am Soc Nephrol* 2004; 15: 2959–2964
- Eifinger F, Wahn F, Quersfeld U *et al.* Coronary artery calcifications in children and young adults treated with renal replacement therapy. *Nephrol Dial Transplant* 2000; 15: 1892–1894
- Briese S, Wiesner S, Will JC *et al.* Arterial and cardiac disease in young adults with childhood-onset end-stage renal disease-impact of calcium and vitamin D therapy. *Nephrol Dial Transplant* 2006; 21: 1906–1914
- Grases F, Santiago C, Simonet BM *et al.* Sialolithiasis: mechanism of calculi formation and etiologic factors. *Clin Chim Acta* 2003; 334: 131–136
- Nicholson JF, Pesce MA. Reference ranges for laboratory tests and procedures laboratory testing in infants and children. In: Behrman RE, Kliegman RA, Jenson HB (eds). *Nelson Textbook of Pediatrics*. Philadelphia, PA: Saunders Elsevier, 2007, 2943–2949

Received for publication: 11.6.08; Accepted in revised form: 19.2.09